

Educational Material for Parents and Students (Content Meets MDCH Requirements)

Sources: Michigan Department of Community Health, CDC and the National Operating Committee Standards for Athletic Equipment (NOCSAE)

UNDERSTANDING CONCUSSION

Some Common Symptoms

Headache
Pressure in the Head
Nausea/Vomiting
Dizziness

Balance Problems
Double Vision
Blurry Vision
Sensitive to Light

Sensitive to Noise
Sluggishness
Haziness
Fogginess
Grogginess

Poor Concentration
Memory Problems
Confusion
“Feeling Down”

Not “Feeling Right”
Feeling Irritable
Slow Reaction Time
Sleep Problems

WHAT IS A CONCUSSION?

A concussion is a type of traumatic brain injury that changes the way the brain normally works. A concussion is caused by a fall, bump, blow, or jolt to the head or body that causes the head and brain to move quickly back and forth. A concussion can be caused by a shaking, spinning or a sudden stopping and starting of the head. Even a “ding,” “getting your bell rung,” or what seems to be a mild bump or blow to the head can be serious. A concussion can happen even if you haven’t been knocked out.

You can’t see a concussion. Signs and symptoms of concussions can show up right after the injury or may not appear or be noticed until days or weeks after the injury. If the student reports any symptoms of a concussion, or if you notice symptoms yourself, seek medical attention right away. A student who may have had a concussion should not return to play on the day of the injury and until a health care professional says they are okay to return to play.

IF YOU SUSPECT A CONCUSSION:

- SEEK MEDICAL ATTENTION RIGHT AWAY** – A health care professional will be able to decide how serious the concussion is and when it is safe for the student to return to regular activities, including sports. Don’t hide it, report it. Ignoring symptoms and trying to “tough it out” often makes it worse.
- KEEP YOUR STUDENT OUT OF PLAY** – Concussions take time to heal. Don’t let the student return to play the day of injury and until a health care professional says it’s okay. A student who returns to play too soon, while the brain is still healing, risks a greater chance of having a second concussion. Young children and teens are more likely to get a concussion and take longer to recover than adults. Repeat or second concussions increase the time it takes to recover and can be very serious. They can cause permanent brain damage, affecting the student for a lifetime. They can be fatal. It is better to miss one game than the whole season.
- TELL THE SCHOOL ABOUT ANY PREVIOUS CONCUSSIONS** – Schools should know if a student had a previous concussion. A student’s school may not know about a concussion received in another sport or activity unless you notify them.

SIGNS OBSERVED BY PARENTS:

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Can’t recall events prior to or after a hit or fall
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes

CONCUSSION DANGER SIGNS

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. A student should receive immediate medical attention if after a bump, blow, or jolt to the head or body s/he exhibits any of the following danger signs:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that gets worse
- Weakness, numbness, or decreased
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people/places coordination
- Becomes increasingly confused, restless or agitated
- Has unusual behavior
- Loses consciousness (even a brief loss of consciousness should be taken seriously.)

HOW TO RESPOND TO A REPORT OF A CONCUSSION:

If a student reports one or more symptoms of a concussion after a bump, blow, or jolt to the head or body, s/he should be kept out of athletic play the day of the injury. The student should only return to play with permission from a health care professional experienced in evaluating for concussion. During recovery, rest is key. Exercising or activities that involve a lot of concentration (such as studying, working on the computer, or playing video games) may cause concussion symptoms to reappear or get worse. Students who return to school after a concussion may need to spend fewer hours at school, take rest breaks, be given extra help and time, spend less time reading, writing or on a computer. After a concussion, returning to sports and school is a gradual process that should be monitored by a health care professional.

Remember: Concussion affects people differently. While most students with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

To learn more, go to www.cdc.gov/concussion.

Concussion

INFORMATION SHEET



This sheet has information to help protect your children or teens from concussion or other serious brain injury. Use this information at your children's or teens' games and practices to learn how to spot a concussion and what to do if a concussion occurs.

What Is a Concussion?

A concussion is a type of traumatic brain injury—or TBI—caused by a bump, blow, or jolt to the head or by a hit to the body that causes the head and brain to move quickly back and forth. This fast movement can cause the brain to bounce around or twist in the skull, creating chemical changes in the brain and sometimes stretching and damaging the brain cells.

How Can I Help Keep My Children or Teens Safe?

Sports are a great way for children and teens to stay healthy and can help them do well in school. To help lower your children's or teens' chances of getting a concussion or other serious brain injury, you should:

- Help create a culture of safety for the team.
 - Work with their coach to teach ways to lower the chances of getting a concussion.
 - Talk with your children or teens about concussion and ask if they have concerns about reporting a concussion. Talk with them about their concerns; emphasize the importance of reporting concussions and taking time to recover from one.
 - Ensure that they follow their coach's rules for safety and the rules of the sport.
 - Tell your children or teens that you expect them to practice good sportsmanship at all times.
- When appropriate for the sport or activity, teach your children or teens that they must wear a helmet to lower the chances of the most serious types of brain or head injury. However, there is no “concussion-proof” helmet. So, even with a helmet, it is important for children and teens to avoid hits to the head.

 **Plan ahead.** What do you want your child or teen to know about concussion?

How Can I Spot a Possible Concussion?

Children and teens who show or report one or more of the signs and symptoms listed below—or simply say they just “don't feel right” after a bump, blow, or jolt to the head or body—may have a concussion or other serious brain injury.

Signs Observed by Parents or Coaches

- Appears dazed or stunned
- Forgets an instruction, is confused about an assignment or position, or is unsure of the game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes
- Can't recall events *prior to* or *after* a hit or fall

Symptoms Reported by Children and Teens

- Headache or “pressure” in head
- Nausea or vomiting
- Balance problems or dizziness, or double or blurry vision
- Bothered by light or noise
- Feeling sluggish, hazy, foggy, or groggy
- Confusion, or concentration or memory problems
- Just not “feeling right,” or “feeling down”

Talk with your children and teens about concussion. Tell them to report their concussion symptoms to you and their coach right away. Some children and teens think concussions aren't serious, or worry that if they report a concussion they will lose their position on the team or look weak. Be sure to remind them that *it's better to miss one game than the whole season.*



CONCUSSIONS AFFECT EACH CHILD AND TEEN DIFFERENTLY.

While most children and teens with a concussion feel better within a couple of weeks, some will have symptoms for months or longer. Talk with your children's or teens' healthcare provider if their concussion symptoms do not go away, or if they get worse after they return to their regular activities.

What Are Some More Serious Danger Signs to Look Out For?

In rare cases, a dangerous collection of blood (hematoma) may form on the brain after a bump, blow, or jolt to the head or body and can squeeze the brain against the skull. Call 9-1-1 or take your child or teen to the emergency department right away if, after a bump, blow, or jolt to the head or body, he or she has one or more of these danger signs:

- One pupil larger than the other
- Drowsiness or inability to wake up
- A headache that gets worse and does not go away
- Slurred speech, weakness, numbness, or decreased coordination
- Repeated vomiting or nausea, convulsions or seizures (shaking or twitching)
- Unusual behavior, increased confusion, restlessness, or agitation
- Loss of consciousness (passed out/knocked out). Even a brief loss of consciousness should be taken seriously

Children and teens who continue to play while having concussion symptoms, or who return to play too soon—while the brain is still healing—have a greater chance of getting another concussion. A repeat concussion that occurs while the brain is still healing from the first injury can be very serious, and can affect a child or teen for a lifetime. It can even be fatal.

What Should I Do If My Child or Teen Has a Possible Concussion?

As a parent, if you think your child or teen may have a concussion, you should:

1. Remove your child or teen from play.
2. Keep your child or teen out of play the day of the injury. Your child or teen should be seen by a healthcare provider and only return to play with permission from a healthcare provider who is experienced in evaluating for concussion.
3. Ask your child's or teen's healthcare provider for written instructions on helping your child or teen return to school. You can give the instructions to your child's or teen's school nurse and teacher(s) and return-to-play instructions to the coach and/or athletic trainer.

Do not try to judge the severity of the injury yourself. Only a healthcare provider should assess a child or teen for a possible concussion. Concussion signs and symptoms often show up soon after the injury. But you may not know how serious the concussion is at first, and some symptoms may not show up for hours or days.

The brain needs time to heal after a concussion. A child's or teen's return to school and sports should be a gradual process that is carefully managed and monitored by a healthcare provider.

To learn more, go to [cdc.gov/HEADSUP](https://www.cdc.gov/HEADSUP)

